

6-Gingerol inhibits hair shaft growth in cultured human hair follicles and modulates hair growth in mice

Yong Miao

Friday, September 4, 2020 11:13 AM

Introduction

This study investigates if ginger affects hair growth.

Conclusions

- Ginger does inhibit hair growth.
- Ginger inhibits hair growth, likely by a mitochondria mediated apoptosis (cell death).
- Ginger inhibits hair growth by inhibiting dermal papillary cells that regulate hair growth.

Amendments

OPEN ACCESS Freely available online



6-Gingerol Inhibits Hair Shaft Growth in Cultured Human Hair Follicles and Modulates Hair Growth in Mice

Yong Miao¹, Yabin Sun², Wenjun Wang³, Benjun Du¹, Shun-e Xiao¹, Yijue Hu¹, Zhiqi Hu^{1*}

1 Department of Plastic and Aesthetic Surgery, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China, **2** GCP Office, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China, **3** Cancer Research Institute, Southern Medical University, Guangzhou, Guangdong, China

Abstract

Ginger (*Zingiber officinale*) has been traditionally used to check hair loss and stimulate hair growth in East Asia. Several companies produce shampoo containing an extract of ginger claimed to have anti-hair loss and hair growth promotion properties. However, there is no scientific evidence to back up these claims. This study was undertaken to measure 6-gingerol, the main active component of ginger, on hair shaft elongation *in vitro* and hair growth *in vivo*, and to investigate its effect on human dermal papilla cells (DPCs) *in vivo* and *in vitro*. 6-Gingerol suppressed hair growth in hair follicles in culture and the proliferation of cultured DPCs. The growth inhibition of DPCs by 6-gingerol *in vitro* may reflect a decrease in the Bcl-2/Bax ratio. Similar results were obtained *in vivo*. The results of this study showed that 6-gingerol does not have the ability to promote hair growth, on the contrary, can suppress human hair growth via its inhibitory and pro-apoptotic effects on DPCs *in vitro*, and can cause prolongation of telogen phase *in vivo*. Thus, 6-gingerol rather than being a hair growth stimulating drug, it is a potential hair growth suppressive drug; i.e. for hair removal.

Citation: Miao Y, Sun Y, Wang W, Du B, Xiao S-e, et al. (2013) 6-Gingerol Inhibits Hair Shaft Growth in Cultured Human Hair Follicles and Modulates Hair Growth in Mice. PLoS ONE 8(2): e57226. doi:10.1371/journal.pone.0057226

Editor: Michael Hamblin, MGHL, MMS, United States of America

Received: July 28, 2012; **Accepted:** January 18, 2013; **Published:** February 21, 2013

Copyright: © 2013 Miao et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Natural Science Foundation of China (Grant No. 31170949). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: doctorhq@hotmail.com

Introduction

Hair is a protective appendage on the body that is considered an accessory structure of the integument. Although the perturbation and loss of hair follicles and alterations in hair fiber production in humans are generally not life-threatening events, their profound impact on social interactions and on patients' psychological well-being is undeniable [1,2] which in turn increases the demands for the possible treatment options and turns hair loss treatment products into a multi-million dollar industry. Despite this, most current products are ineffective, as evidenced by the fact that the FDA has approved only two treatments for hair loss [3]. Given the limited, transient and somewhat unpredictable efficacy of the current approved anti-hair loss medicines [3], novel pharmacological treatments and agents are on demand. There have been many attempts to discover materials from traditional herbal medicines that prevent hair loss [4–6]. For instance, green tea, the tuber flower (*Polygonum multiflorum* Thunb) and ginger (*Zingiber officinale*) have been traditionally used in East Asia to treat patients suffering from baldness and hair loss. Several companies produce shampoos containing an extract of *P. multiflorum* or ginger extracts which are claimed to have anti-hair loss and hair growth promotion properties [9,10]. Recently, several studies demonstrated that the green tea [4] or *P. multiflorum* extract [8] can promote hair growth; however, there is no scientific evidence to support this claim for ginger.

The mammalian hair follicle contains dermal papilla (DP), which mainly consist of DP cells (DPCs), and dermal sheath derived from the mesenchyme as well as epithelial cells of the outer root sheath, inner root sheath, matrix and hair shaft, derived from

the epithelium [11]. The postnatal hair follicle undergoes a cycle of anagen (growth phase), catagen (regression phase) and telogen (resting phase). The reciprocal interactions between the epithelium and mesenchyme are essential for postnatal hair growth and cycling of hair follicles. The DP is known to have a key role in the regulation of hair growth and cycling [11]. Thus, any factor that affects the functions of DPCs can influence hair growth. For instance, Minoxidil [12] and epigallocatechin-3-gallate [4] stimulate the hair growth by exerting their anti-apoptotic effects on DPCs (through increasing the Bcl-2/Bax ratio), this is one of the mechanism of actions on Minoxidil. Cisplatin [13] on the other hands leads to hair loss by inducing apoptotic effects on DPCs (through decreasing the Bcl-2/Bax ratio).

There are several recent scientific investigations aimed at the isolation and identification of active constituents of ginger and the scientific verification of its pharmacological actions and those of its constituents [14,15]. The results show that 6-gingerol is the most abundant active constituent and has various pharmacological and microbiological effects, including anti-tumorigenic, anti-inflammatory, anti-oxidant and anti-emetic actions; However, pharmacological effects of 6-gingerol on hair growth are not scientifically proven.

In this study, we have investigated the effects of 6-gingerol on human DPCs and hair shaft growth *ex vivo* and hair growth in mice.

Materials and Methods

2.1. Isolation and culture of human hair follicles

Punch biopsy (4 mm) specimens were taken from male nonbalding occipital scalps of patients undergoing hair transplan-

The hair follicle is a pocket that extends lengthwise upward to create a "cave" where the hair shaft is created by matrix cells (also known as epithelial germination cells) started at the hair bulb. These cells proliferate (multiply) and differentiate (turn into) into keratin producing cells (keratinocytes), which create keratin protein into hair fibers, creating the hair shaft, while also integrating into the hair shaft and killing themselves, creating keratinized cells. This whole process is controlled by dermal papillary cells (DPCs) that send signals to the matrix cells to divide/multiply/proliferate and differentiate to migrate upward from the hair bulb and into the hair shaft to be keratinized cells.






Figure 1. Growth inhibition and induction of apoptosis effects of 6-gingerol on human DPCs *in vitro*. DPCs were cultured with different concentrations of 6-gingerol for 48 h. Compared to the vehicle-treated control, treatment with 10 µg/ml 6-gingerol significantly inhibited the proliferation of cells. (A) Cell viability was determined by the MTT assay. * $P < 0.05$ vs the vehicle-treated control. (B) Apoptosis analysis of cells. doi:10.1371/journal.pone.0057226.g001

tation surgery for androgenic alopecia. The Medical Ethical Committee of the Southern Medical University approved all studies described here. The study was conducted according to the Declaration of Helsinki Principles and informed written consent was obtained from all patients. Hair follicles were isolated and cultured as described by Philpott et al [16]. Briefly, hair follicles were isolated with forceps under a binocular light microscope and cultured in 24-well dishes for 7 days in Williams E medium (Gibco BRL, Gaithersburg, MD) supplemented with 10 ng/ml hydrocortisone, 10 mg/ml insulin, 2 mM L-glutamine, 100 U/ml streptomycin and 10 mg/ml hydrocortisone at 37 °C in a 5% (v/v) CO₂ atmosphere. 6-Gingerol (>95% purity; Sigma Chemical Co., St. Louis, MO) was dissolved in dimethylsulfoxide (DMSO; Solarbio Science & Technology Co., Ltd., Beijing, China) and to keep final concentration of the vehicle < 0.1%. 6-Gingerol was added to culture medium at 10 or 20 µg/ml, and 0.1% (v/v) DMSO was used as a control. In all experiments, culture medium and 6-gingerol were refreshed every other day. A total of 100 anagen hair follicles from 3 different volunteers (60 follicles/subject) were cultured under each growth condition. The mean length of hair follicle from the bottom of the dermal papillae at day zero was 4.5 mm.

2.2. Culture of human DPCs

DPCs were isolated and cultured as described by Magrel et al [17]. Briefly, dermal papillae were microdissected from the bulbs of dissected hair follicles, transferred onto plastic dishes and cultured in Dulbecco's modified Eagle medium (DMEM; Gibco, Grand Island, NY) supplemented with 100 U/ml penicillin, 100 mg/ml streptomycin and 20% (v/v) fetal bovine serum (FBS; Gibco) at 37 °C in a humidified 5% CO₂ atmosphere. The explants were kept in the medium for 7 days and the medium was changed every 3 days. Once cell outgrowth was sub-confluent, cells were harvested with 0.25% (w/v) trypsin-EDTA (Invitrogen, Carlsbad, CA) and subcultured with a split ratio of 1:3. Afterwards, DPCs were maintained in DMEM supplemented with 10% FBS.

2.3 Cell proliferation and apoptosis assays

Cell proliferation was determined using the MTT assay as described by Mosmann [18]. Briefly, DPCs (3×10^4 cells/well) were seeded into 96-well plates and incubated for 24 h before adding 6-gingerol at 5 or 10 µg/ml and then incubated at 37 °C for 48 h, using 0.1% DMSO as a negative control. Absorbance at 490 nm was measured using an ELISA reader. Relative cell viability was calculated by dividing the absorbance of treated cells by that of the nontreated cells in each experiment.




Figure 2. Effect of 6-gingerol on the expression of Bcl-2 and Bax proteins in cultured human DPCs. (A) DPCs were treated with the indicated concentrations of 6-gingerol (0.1% DMSO control) for 48 h and subjected to western blot analysis. Representative bands are shown. (B) Histograms showing the quantification of western blot bands. GAPDH was used as the internal control. * $P < 0.05$ compared to the control (0.1% DMSO) group. doi:10.1371/journal.pone.0057226.g002

Figure 1

This data shows dermal papillary cells (see introduction for details) that control the growth of the hair follicle being plated onto dishes and [1A] ginger is being added to the plate in increasing concentrations to determine cell growth/multiplying, while [1B] is a determination of apoptosis (cell death), with more black dots (representative of cells) located higher up being an indication of more cell death (apoptosis).

Primary Results

- There is a decrease in dermal papillary cell (DPC) growth with the addition of ginger.
- There is an increase in DPC death with the addition of ginger.

Take Away: In isolated cells, ginger inhibits the growth of hair growth regulating cells.

Figure 2

[2A] This data here is showing the amount of Bax (pro-apoptosis/cell death) protein and BCL-2 (anti-apoptosis/cell death) protein levels (darker the smudge, the more there is) in the absence of presence of ginger in PDCs (papillary dermal cells). [2B] This is the graph version quantifying what is seen in 2A.

Primary Results

- Bax increases with ginger addition.
- BCL decreases with ginger addition.

Take Away: This data indicates there is an increase in pro-apoptosis (cell death) signaling and a reduction in anti-apoptosis signaling.




Figure 3. Growth inhibitory effect of 6-gingerol on the elongation of human hair shafts *in vitro*. Hair follicles were treated with different concentrations of 6-gingerol for 7 days and the elongation of human hair shafts was measured using a calibrated eyepiece graticule in a light microscope at a magnification of 20 \times . (n = 60)
doi:10.1371/journal.pone.0057226.g003

DPCs seeded in 6-well plates at a density of 2.5×10^3 cells/well were treated with 5 or 10 $\mu\text{g/ml}$ 6-gingerol or 0.1% DMSO as a control for 48 h. Cell apoptosis was examined using an Annexin V detection kit (Caltag-Mediatech, Buckingham, UK) according to the manufacturer's instructions. Data acquisition and analysis were done in a FACSort Cytometer (FAC-SCA, New York, NY). A minimum of 2.5×10^3 cell events was acquired for each group in the process of analysis and each experiment was repeated at least 3 times.

2.4 Western blot analysis

Both control and 6-gingerol-treated DPCs were collected at 48 h after treatment. Total cell lysates were prepared and 30 μg of protein was subjected to sodium dodecyl sulfate/polyacrylamide gel electrophoresis (SDS-PAGE) followed by immunoblot analysis. Primary antibodies were incubated at appropriate dilutions: anti-Bcl-2 monoclonal antibody, 1:500; anti-Bax monoclonal antibody, 1:500; and anti-GAPDH monoclonal antibody, 1:1000 (Santa Cruz Biotechnology Inc., Santa Cruz, CA). The immune complexes were detected using a western blotting enhanced chemiluminescence (ECL) kit (Santa Cruz) and quantified using analysis/PC densitometry software (Bio-Rad Laboratories, Hercules, CA).

Table 1. Elongation of hair shafts with the indicated treatment (n = 60).

Treatment	Hair shaft length (mm)	*P
Control (0.1% v/v DMSO)	1.48 \pm 0.20	
6-Gingerol (10 $\mu\text{g/ml}$)	1.41 \pm 0.23	0.11
6-Gingerol (20 $\mu\text{g/ml}$)	1.06 \pm 0.21	0.00*

*Versus the control.
doi:10.1371/journal.pone.0057226.t001

2.5 In vivo analysis of 6-gingerol

A total of 36 female C57BL/6 mice (8 weeks old) were purchased from the Experimental Animal Center of Southern Medical University (Guangzhou, China) and divided randomly into 3 groups (12/group). Anagen was induced by depilation of skin on the back of C57BL/6 mice that were in the telogen phase of the cycle, as described [19]. Briefly, hair was removed by topical application of calcium thioglycolate. From the following day (day 1), 0.2 ml of 0.1 mg/ml or 1 mg/ml 6-gingerol in 75% (v/v) ethanol was applied to the dorsal skin of each animal every day for 10 days. Control animals received the vehicle solution alone. The back of each mouse was observed and photographed every 5 days following depilation. All experiments lasted for 20 days and the mice were then sacrificed. The histology of skin specimens taken from the back of each mouse was examined with hematoxylin and eosin (H&E) staining. All experimental manipulations were done in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and with the approval of the Experimental Animal Ethical Committee of Southern Medical University.

2.6 Statistical analysis

All data are expressed as mean \pm SD from three independent experiments. All statistical analysis was done with SPSS 13.0. Differences between experimental groups were evaluated by Student's t-test. Statistically significant difference was set at $P < 0.05$.

Results

3.1 6-Gingerol significantly inhibits the growth of DPCs and induces cell apoptosis

To examine the effect of 6-gingerol on the proliferation of DPCs, we treated DPCs with DMSO (control) or 6-gingerol (5 or 10 $\mu\text{g/ml}$). Within 48 h of culture, 6-gingerol exhibited a significant dose-dependent growth inhibitory effect on DPCs (Fig. 1A). To determine the mechanisms underlying 6-gingerol-

Figure 3 & Table 1

Here, the researchers are showing images of isolated hair shafts (the hair itself) surrounded by their hair follicle (the hair environment) with increasing concentrations of ginger exposed. Table 1 is a quantification of growth of figure 3.

Primary Results

- Ginger addition trends toward a significant hair inhibitory effect with 10 $\mu\text{g/ml}$ concentration of ginger.
- Ginger addition at double the concentration (20 $\mu\text{g/ml}$) leads to a clear hair growth inhibition.

Take Away: This shows that ginger inhibits hair growth when the shaft and the entire follicle are present.



Figure 4. Inhibition of anagen induction from telogen by 6-gingerol in C57BL/6 mice. After depilation, the skin on the back was treated with vehicle (right) or 1 mg/ml 6-gingerol (left) every day for 10 days. Photographs of each animal were taken every 5 days. Compared to the vehicle-treated control, 1 mg/ml 6-gingerol can significantly inhibit anagen induction from telogen. (A) Day zero, (B) 5 days, (C) 10 days, (D) 15 days and (E) 20 days after depilation. (F) Enlarged photograph of the left-hand mouse in (E). doi:10.1371/journal.pone.0057226.g004

inhibited DPCs growth, the effect of 6-gingerol on cell apoptosis was assessed by staining with fluorescein isothiocyanate (FITC) and Annexin V. Consistently, the exposure of DPCs to 6-gingerol significantly increased cell apoptosis as compared to the control group, suggesting apoptosis as the mechanism underlying 6-gingerol-induced cytotoxicity (Fig. 1B).

3.2 Effects of treatment with 6-gingerol on apoptosis regulatory proteins

Earlier, 6-gingerol was shown to induce apoptosis through the mitochondrial death pathway [20–22]. Because this pathway of apoptosis is known to be regulated by the balance of anti- and pro-apoptotic proteins in the Bcl-2 family [23,24], we examined the level of expression of two key Bcl-2 family proteins Bcl-2 and Bax in response to treatment of DPCs with 6-gingerol. Immunoblotting studies show that 6-gingerol induced significant down-regulation of the anti-apoptotic Bcl-2 protein and up-regulation of the

apoptotic Bax protein compared with the vehicle-treated control. Additionally, Bax protein was up-regulated in a 6-gingerol dose-dependent manner ($P < 0.05$) (Fig. 2A, B).

3.3 6-Gingerol inhibits hair shaft elongation in cultured human hair follicles

Because DP has a key role in the regulation of hair growth, we investigated the effect of 6-gingerol on hair shaft elongation. Human scalp hair follicles were isolated and cultured both in the absence and in the presence of 6-gingerol. We observed that 20 μ g/ml 6-gingerol significantly inhibits elongation of hair shafts (Fig. 3 and Table 1).

3.4 6-Gingerol delays anagen induction from telogen in mice

The results given above show only the effect of 6-gingerol in cultured DPCs and hair follicles *in vitro*, which is not the same as the situation *in situ*. To overcome this limitation, we assessed the role of 6-gingerol in hair growth *in vivo* using 8 weeks old C57BL/6 mice. When the dorsal pelage hair follicles were in the telogen phase, mice were shaved on the back and those showing a uniform telogen stage skin were chosen. The dorsal skin was treated with vehicle alone (control) or with 6-gingerol every day for 10 days. Figure 4 shows hair regrowth at 5-day intervals for the control and for treatment with 6-gingerol (1 mg/ml 6-gingerol as an example). In normal control mice, faint hair regrowth was observed at 10 days after depilation. It became evident that hair on the mice had fully regrown to the original status within ~20 days. After the topical application of 1 mg/ml 6-gingerol, faint hair regrowth was observed at 15 days after depilation and sparse hair regrowth was observed at 20 days after depilation. The effect of 6-gingerol on the density of hair follicles was further assessed by staining with H&E. Consistently, topical application of 6-gingerol can significantly decrease the number of hair follicles as compared to the control group (Fig. 5).

Discussion

Ginger (*Z. officinale*) has been traditionally used in East Asia to prevent hair loss and stimulate hair growth. Currently, several companies produce shampoos containing ginger extract and claim




Figure 5. 6-Gingerol decreased the number of hair follicles in C57BL/6 mice. After depilation the back, the skin was treated with vehicle alone (control) or 1 mg/ml 6-gingerol every day for 10 days. The effect of 6-gingerol on the hair follicles was analyzed using staining with H&E at 20 days after depilation. (a)–(c) Longitudinal sections of the dorsal skins. (b)–(d) Transverse sections of the dorsal skins. Each scale bar represents 200 μ m. doi:10.1371/journal.pone.0057226.g005

Figure 4

These images are mice that have had their backs shaved the same way and ginger is added to one of the mice's back skin (left mouse) and the other is exposed to a negative control (right mouse).

Primary Results

- Mouse hair growth is greater in the right mouse, especially evident in panel D.

Take Away: This shows a "live system" that ginger inhibits hair growth.

that it has anti-hair loss and hair growth promoting effects [9,10]; however, there is no convincing evidence to support the claimed effects of ginger on hair growth. In this study, 6-gingerol, the main active component of ginger, was found to significantly inhibit hair growth both *in vivo* and *in vitro*. To our knowledge, the present study is the first to evaluate the effect of 6-gingerol on hair growth. Our results show that 6-gingerol does not promote hair growth and can inhibit hair growth by inducing apoptotic effects on DPCs (by decreasing the Bcl-2/Bax ratio).

We found that 6-gingerol in the concentration range 5–10 µg/ml caused a significant dose-dependent inhibition of the proliferation of human DPCs *ex vivo*. Moreover, 20 µg/ml 6-gingerol significantly suppressed elongation of the hair shaft (1.06 ± 0.21 mm) compared to the control group (1.48 ± 0.20 mm).

6-Gingerol has been shown to induce apoptosis mainly through stimulation of mitochondrial (intrinsic) death pathway [20–22], which is regulated primarily by Bcl-2 family proteins, notably the pro-apoptotic Bax and anti-apoptotic Bcl-2 proteins [23,24]. It is generally accepted that Bcl-2 protects cells from apoptosis and that the activity of Bcl-2 is determined by interaction with Bax [25]. Interestingly, Bcl-2 and Bax have been shown to regulate hair follicle apoptosis during the apoptosis-driven regression phase of the hair cycle (catagen) [26,27]. These findings have led to the hypothesis that 6-gingerol induces apoptosis in hair follicle cells by regulating the levels of Bcl-2 and Bax. In this study, we showed that 6-gingerol affects the expression of Bcl-2 and Bax in cultured human DPCs. We found 6-gingerol increased Bax and decreased Bcl-2 expression in cultured DPCs. Treatment with 6-gingerol for 48 h induced a significant increase in the expression of Bax and decrease in the expression of Bcl-2 compared with the control group. Additionally, Bax protein was up-regulated in a 6-gingerol dose-dependent manner. On the basis of the results of this and earlier research [4,12,13,26,27], we propose that 6-gingerol induces apoptosis in DPCs by decreasing the level of Bcl-2 and increasing the level of Bax.

We have confirmed that 6-gingerol suppresses hair shaft elongation *ex vivo* by inducing apoptotic damage in DPCs, but it is not clear whether 6-gingerol can penetrate into hair follicles and cause apoptotic damage in DPCs *in vivo*. Compared to the situation *ex vivo*, DPCs in hair follicles *in situ* have a higher resistance to damage factors [28]; thus, to determine whether the changes observed *ex vivo* occurred *in vivo*, we investigated the hair growth-inhibiting activity of 6-gingerol using 8 weeks old C57BL/6 mice in the stable and consistent telogen phase [19]. The shaved

back skin of C57BL/6 mice was treated with daily topical application of 6-gingerol for 10 days. In normal control mice, faint hair regrowth was observed at 10 days and full regrowth was observed at 20 days after depilation. However, in the 1 mg/ml 6-gingerol group, only faint hair regrowth was observed at 15 days and sparse regrowth was observed at 20 days after depilation. To further investigate the hair growth inhibition, we assessed the effect of 6-gingerol on the density of hair follicles by staining with (H&E). Consistently, topical application of 1 mg/ml 6-gingerol can substantially decrease the number of hair follicle as compared to the control group. Thus, it was confirmed that the events initially observed *in vitro* actually occurred *in vivo*.

Unwanted body hair can be emotionally and socially devastating [29], resulting in the search for various treatment modalities. The simplest and most popular method is the use of depilatory materials, which chemically remove hair from the skin surface by dissolving keratin but have only a temporary effect. Unfortunately, there are some side-effects, such as skin allergy, parafollicular hyperkeratosis or skin burns [30]. According to the results of this present study, 6-gingerol can decrease the density of hair follicles via inducing apoptotic effects on DPCs, and is a potential permanent hair removal drug.

In summary, we report for the first time that 6-gingerol has no effect on promoting hair growth, on the contrary, can suppress human hair growth via its inhibitory and pro-apoptotic effects on DPCs *in vitro*, and cause prolongation of telogen phase *in vivo*. Thus, 6-gingerol rather than being a hair growth stimulating drug, it is a potential hair growth suppressive drug, i.e. for hair removal. The inadequacies of this study are that this research has slightly fewer cases, the unknown effect of 6-gingerol on follicular epithelial cells and genetic changes of 6-gingerol on dermal papilla cells *in vivo*. Therefore, in future experiments, besides increasing the sample size, we should do further experiments to address the above-mentioned problems.

Acknowledgments

The authors thank Shengji Wang, Boxin Zhao, Jindou Jiang, Zhenhua Li and Zhidan Zhang for their technical support and critical discussions.

Author Contributions

Conceived and designed the experiments: ZH YM. Performed the experiments: YM YS SX. Analyzed the data: WW BJ. Contributed reagents/materials/analysis tools: YH YS. Wrote the paper: YM.

References

- Cash TF. (1999) The psychosocial consequences of androgenetic alopecia: a review of the research literature. *Br J Dermatol* 141:396–405.
- Schmidt S, Fischer TW, Chen MM, Strauss BM, Elner F. (2001) Strategies of coping and quality of life in women with alopecia. *Br J Dermatol* 146:1038–1043.
- Prior VH. (1999) Treatment of hair loss. *N Engl J Med* 341:964–973.
- Kwon OK, Han JH, Yoo HG, Chung JH, Cho KH, et al. (2005) Human hair growth enhancement *in vitro* by green tea epigallocatechin-3-gallate (EGCG). *Phytotherapy* 14:551–555.
- Kawano M, Han J, Kikuchi M, Ioda H. (2005) Hair growth regulation by the extract of aromatic plant *Erica multiflora*. *J Nat Med* 63:335–339.
- Datta K, Singh AT, Mukherjee A, Bhat B, Ramesh B, et al. (2009) Eclipta alba extract with potential for hair growth promoting activity. *J Ethnopharmacol* 124:450–456.
- Kim SC, Kang JI, Kim MK, Hyan JH, Boo HJ, et al. (2010) Promotion effect of monggolantennin, a component of Citrus aurantium, on hair growth. *Eur J Dermatol* 20:47–50.
- Park HJ, Zhang N, Park DK. (2011) Topical application of Polygonum multiflorum extract induces hair growth of resting hair follicles through upregulating Shh and β-catenin expression in C57BL/6 mice. *J Ethnopharmacol* 133:369–373.
- Alfaba website. Available: http://www.alfaba.com/products/236088744/ginger_shampoo.html
- DHgate website. Available: <http://factory.dhgate.com/mens-hair-care-products/ginger-hair-loss-prevention-hair-shampoo-p4962195.html>
- Conrad G, Paus R. (1999) The biology of hair follicles. *N Engl J Med* 341:491–497.
- Han JH, Kwon OS, Chung JH, Cho KH, Eun HC, et al. (2004) Effect of minoxidil on proliferation and apoptosis in dermal papilla cells of human hair follicle. *J Dermatol Sci* 34:91–98.
- Luangpiang S, Nimmamanit U, Charochoke P, Leonard SS, Pongkhamanon V, et al. (2011) Hydroxyl radical mediates cisplatin-induced apoptosis in human hair follicle dermal papilla cells and keratinocytes through Bcl-2-dependent mechanism. *Apoptosis* 16:709–722.
- Alf B, Bhandari G, Tanira MO, Nemmar A. (2008) Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food Chem Toxicol* 46:609–620.
- Bağcı MS, Haniyazlar R, Pereira MM, D'Souza JJ, Palfay PL, et al. (2011) Update on the chemopreventive effects of ginger and its phytochemicals. *Crit Rev Food Sci Nutr* 51:409–523.
- Philpott MP, Green MR, Krulic T. (1990) Human hair growth *in vitro*. *Cell Sci* 97:471–476.
- Magot M, Kanar S, Paus R, Tabin DJ. (2002) Simple and rapid method to isolate and culture follicular papillae from human scalp hair follicles. *Exp Dermatol* 11:381–385.

18. Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65:55–63.
19. Miller-Rosov S, Handjiski B, Eichmüller S, Fritsch K, M-Kay IA, et al. (2001) A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J Invest Dermatol* 117:3–15.
20. Shukla Y, Prasad S, Tripathi C, Singh M, George J. (2007) In vitro and in vivo modulation of testosterone mediated alterations in apoptosis related proteins by [6]-gingerol. *Mol Nutr Food Res* 51:1492–1502.
21. Nagam N, George J, Srivastava S, Rao P, Bhal K, et al. (2010) Induction of apoptosis by [6]-gingerol associated with the modulation of p53 and involvement of mitochondrial signaling pathway in B[a]P-induced mouse skin tumorigenesis. *Cancer Chemother Pharmacol* 65:987–996.
22. Lee C, Park GH, Kim CY, Jung JH. (2011) [6]-Gingerol attenuates β -amyloid-induced oxidative cell death via fortifying cellular antioxidant defense system. *Food Chem Toxicol* 49:1261–1269.
23. Mignone B, Vayssiere JC. (1996) Mitochondria and apoptosis. *Eur J Biochem* 232:1–15.
24. Tsujimoto Y. (1998) Role of Bcl-2 family proteins in apoptosis: apoptosomes or mitochondria. *Genes Cells* 3:697–707.
25. Adams JM, Cory S. (1998) The Bcl-2 protein family: arbiters of cell survival. *Science* 281:1322–1326.
26. Muller-Rosov S, Rosatier H, Lindeker G, Proves EMJ, Kopper TS, et al. (1999) Hair follicle apoptosis and Bcl-2. *J Invest Dermatol Symp Proc* 4:272–277.
27. Botchkareva NV, Ahluwalia G, Shander D (2006) Apoptosis in the hair follicle. *J Invest Dermatol* 126:259–264.
28. Lindeker G, Botchkareva VA, Botchkareva NV, Ling G, van der Veen C, et al. (1997) Analysis of apoptosis during hair follicle regression (catagen). *Am J Pathol* 151:1601–1617.
29. Blume-Peytavi U. (2011) An overview of unexplained female hair. *Br J Dermatol* 165:19–23.
30. Zoumaran J, Kwei JN, Vandervoort J. (2008). A case review of patients presenting to Royal North Shore Hospital, with hair removal wax burns between January and November 2006. *Burns* 34:254–256.